



# Australian Vaccination Schedule

## 1 May 2009

Age	Vaccine	Brands supplied by WA DoH	Disease Prevented
Birth	Hep B	H-B-Vax 11 (Paed)	Hepatitis B
2 months	7vPCV ORV <sup>1</sup> DTPa-IPV-HepB-Hib	Prevenar RotaTeg Infanrix Hexa	Pneumococcal Rotavirus Diphtheria, Tetanus, Pertussis, Poliomyelitis, Hepatitis B and <i>Haemophilus Influenza</i> type b
4 months	7vPCV ORV <sup>1</sup> DTPa-IPV-HepB-Hib	Prevenar RotaTeg Infanrix Hexa	Pneumococcal Rotavirus Diphtheria, Tetanus, Pertussis, Poliomyelitis, Hepatitis B and <i>Haemophilus Influenza</i> type b
6 months	7vPCV ORV <sup>1</sup> DTPa-IPV-HepB-Hib	Prevenar RotaTeg Infanrix Hexa	Pneumococcal Rotavirus Diphtheria, Tetanus, Pertussis, Poliomyelitis, Hepatitis B and <i>Haemophilus Influenza</i> type b
6 months to <5years	Flu (annual) <sup>2</sup>	Fluvax, Vaxigrip	Influenza
12 months	MMR Hib MenCCV	Priorix Hiberix Neisvac-C	Measles, Mumps, Rubella <i>Haemophilus Influenza</i> type b Meningococcal C
<u>Special Populations</u> a. Aboriginal children b. Medically at risk children	Hep A 7vPCV	Vaqa Prevenar	Hepatitis A Pneumococcal
18 months	VZV	Varilrix	Varicella (chickenpox)
<u>Special Populations</u> Aboriginal children	HepA 23vPPV	Vaqa Pneumovax 23	Hepatitis A Pneumococcal
4 years	MMR DTPa-IPV	Priorix Quadracel	Measles, Mumps, Rubella Diphtheria, Tetanus, Pertussis, Poliomyelitis
<u>Special Populations</u> Medically at risk children	23vPPV <sup>3</sup>	Pneumovax 23	Pneumococcal
School Year 7	Hep B (two doses - 4 to 6 months apart) dTpa VZV	H-B-Vax 11 (Adult)  Boostrix or Adacel Varilrix	Hepatitis B  Diphtheria, Tetanus, Pertussis Varicella (chickenpox)
<u>Special Populations</u> Females only	HPV <sup>4</sup>	Gardasil	Human Papilloma Virus
<b>Adults</b> 15-49 yrs Aboriginal with medically risk OR ≥50 years Aboriginal OR Non-Aboriginal ≥ 65 years	23vPPV (two doses - 5 or more years apart)  Flu (annual)	Pneumovax  Fluvax, Vaxigrip	Pneumococcal  Influenza
<u>Special Populations</u> Females 12-26 Years	HPV <sup>4</sup>	Gardasil	Human Papilloma Virus

Annual Influenza vaccine for children aged 6 months to < 5 years

<sup>1</sup> **Rotavirus vaccine (RotaTeg-ORV):** The RotaTeg vaccine is for ORAL use. The first dose of RotaTeg is given at/before 12 weeks of age. The second dose should be given preferably by 28 weeks of age to allow for a minimum interval of 4 weeks before the third dose. The third dose should be given by 32 weeks of age. Minimum interval of 4 weeks. **Late or catch up doses are not to be given.**

<sup>2</sup> **Influenza vaccine for children:** Children who have had 1 or no doses of influenza vaccine ever in their lives should receive 2 doses of vaccine given at least 1 month apart. Children who have had 2 or more doses of Influenza vaccine at any time in their lives prior to the current year, require only 1 dose of vaccine annually.

<sup>3</sup> **Additional Pneumococcal vaccine for medical at risk children:** A 4th dose of 7vPCV at 12 months of age, and a booster dose of 23vPPV at 4 years of age.

<sup>4</sup> **Gardasil 12-26 years females HPV catch up completion date December 2009:** When possible the 3 doses of HPV should be given as per the recommended schedule of 0, 2 and 6 months. First dose needs to be given prior to 30 June. If an accelerated schedule is unavoidable, the second dose should be given at least 1 month after the first dose and 3 months after the second dose.

**IPV (IPOL) for special population groups, e.g. refugee catch-up immunisations:** IPOL can be given to persons needing immunity to polio for whom combination vaccines that contain polio antigens are not indicated. Note: IPOL is not funded as a travel vaccine.

**Groups with special vaccination requirements, e.g. medically at risk:** Refer to the 9th edition Immunisation Handbook pages 75 - 102 for the vaccination guidelines