

West Australian Vaccination 1st April 2010 Schedule

Age	Vaccine	Brands supplied by WA DoH	Disease Prevented
Birth	Hep B	H-B-Vax 11 (Paed)	Hepatitis B
2 months	7vPCV ORV¹ DTPa-IPV-HepB-Hib	Prevenar RotaTeq Infanrix Hexa	Pneumococcal Rotavirus Diphtheria, Tetanus, Pertussis, Poliomyelitis, Hepatitis B and <i>Haemophilus Influenza</i> type b
4 months	7vPCV ORV¹ DTPa-IPV-HepB-Hib	Prevenar RotaTeq Infanrix Hexa	Pneumococcal Rotavirus Diphtheria, Tetanus, Pertussis, Poliomyelitis, Hepatitis B and <i>Haemophilus Influenza</i> type b
6 months	7vPCV ORV¹ DTPa-IPV-HepB-Hib	Prevenar RotaTeq Infanrix Hexa	Pneumococcal Rotavirus Diphtheria, Tetanus, Pertussis, Poliomyelitis, Hepatitis B and <i>Haemophilus Influenza</i> type b
6 months to <5years	Flu (annual)²	Fluvax, Vaxigrip	Influenza
12 months	MMR Hib MenCCV	Priorix Hiberix Neisvac-C	Measles, Mumps, Rubella <i>Haemophilus Influenza</i> type b Meningococcal C
<u>Special Populations</u> a. Aboriginal children b. Medically at risk children	Hep A 7vPCV	Vaqa Prevenar	Hepatitis A Pneumococcal
18 months	VZV	Varilrix	Varicella (chickenpox)
<u>Special Populations</u> Aboriginal children	HepA 23vPPV	Vaqa Pneumovax 23	Hepatitis A Pneumococcal
4 years	MMR DTPa-IPV	Priorix Quadracel	Measles, Mumps, Rubella Diphtheria, Tetanus, Pertussis, Poliomyelitis
<u>Special Populations</u> Medically at risk children	23vPPV³	Pneumovax 23	Pneumococcal
School Year 7	Hep B (two doses – 4 to 6 months apart) dTpa VZV	H-B-Vax 11 (Adult) Boostrix or Adacel Varilrix	Hepatitis B Diphtheria, Tetanus, Pertussis Varicella (chickenpox)
<u>Special Populations</u> Females only	HPV⁴	Gardasil	Human Papilloma Virus
Adults ≥15 yrs Aboriginal OR Non-Aboriginal ≥ 65 years Non-Aboriginal ≥ 65 years, Aboriginal ≥ 50 years, OR Aboriginal ≥15 yrs with a medical at risk condition	Flu (annual)⁵ 23vPPV (two doses – 5 or more years apart)	Fluvax, Vaxigrip, Fluvac Pneumovax	Influenza Pneumococcal
<u>Special Populations</u> Females 12-26 Years	HPV⁴	Gardasil	Human Papilloma Virus

Annual Influenza vaccine for children aged 6 months to < 5 years

¹ **Rotavirus vaccine (RotaTeq-ORV):** The RotaTeq vaccine is for ORAL use. The first dose of RotaTeq is given at/before 12 weeks of age. The second dose should be given preferably by 28 weeks of age to allow for a minimum interval of 4 weeks before the third dose. The third dose should be given by 32 weeks of age. Minimum interval of 4 weeks. **Late or catch up doses are not to be given.**

² **Influenza vaccine for children:** Children who have had 1 or no doses of influenza vaccine ever in their lives should receive 2 doses of vaccine given at least 1 month apart. Children who have had 2 or more doses of Influenza vaccine at any time in their lives prior to the current year, require only 1 dose of vaccine annually.

³ **Additional Pneumococcal vaccine for medical at risk children:** A 4th dose of 7vPCV at 12 months of age, and a booster dose of 23vPPV at 4 years of age.

⁴ **Gardasil 12-26 years females HPV catch up**

⁵ **Influenza vaccine eligibility includes:** All pregnant women, all person 65 years of age or older, all Indigenous Australians aged 15 years and older, all individuals aged six (6) months and over with medical conditions predisposing to severe influenza, namely:

- cardiac disease
- chronic respiratory conditions
- chronic illnesses requiring regular medical follow-up or hospitalisation in the preceding year
- chronic neurological conditions
- people with impaired immunity; and
- children aged six (6) months to 10 years receiving long term aspirin therapy.

IPV (IPOL) for special population groups, e.g. refugee catch-up immunisations: IPOL can be given to persons needing immunity to polio for whom combination vaccines that contain polio antigens are not indicated. Note: IPOL is not funded as a travel vaccine.

Groups with special vaccination requirements, e.g. medically at risk: Refer to the 9th edition Immunisation Handbook pages 75 - 102 for the vaccination guidelines